



The Role of *Olea Europaea* L. Fruit on A2780, A172 and HFFF2 Proliferation

Hadis Bahramiun ^a, Mohammad Kamalinejad ^b, Fatemeh Elahian ^c, Maryam Noubarani ^a, Fatemeh Balaghi ^a,
Fariborz Keyhanfar ^d, Mohammad Reza Eskandari ^a, Farshad H. Shirazi ^{b,e,*}

^a Department of Pharmacology and Toxicology, School of Pharmacy, Zanjan University of Medical Sciences,
Zanjan, Iran.

^b Faculty of Pharmacy, Shahid Beheshti University of Medical Sciences, Tehran. Iran.

^c Cellular and molecular research center, Shahrekord University of Medical Sciences, Shahrekord, Iran.

^d Department of Pharmacology, Faculty of Medicine, Iran University of Medical Sciences, Tehran, Iran

^e SBMU Pharmaceutical Research Center, Tehran, Iran,

Abstract

Olea europaea L. commonly known as olive has been traditionally used for the prevention and treatment of many diseases since ancient times. Olive has been reported to possess a broad spectrum of pharmacological properties. In the present study, we investigated the activity of aqueous extract of *Olea europaea* L. fruit at various concentrations on A2780, A172, and HFFF2 cell lines proliferation by MTT assay. Aqueous extract of olive significantly increased cell proliferation in a dose dependent manner in the cell lines. It has been previously reported that olive has chemopreventive and anti-tumor effects. These disagreements can be explained by differences in cell line properties, type of olive and different solvents in the extracts. However, further investigation is needed to clarify the exact role of olive in cell proliferation and cancer. In this study fruit extract of *Olea europaea* L. showed more activatory effects on A2780 cell line in comparison with A172 and HFFF2. These differences in the activatory effects may be related to the activation of different signaling pathways in different cell lines.

Keywords: Activatory effects, A2780, A172, Cell Proliferation, HFFF2, *Olea europaea* L, Olive.

1. Introduction

Olea europaea L. (Oleaceae) known as olive is native to the Mediterranean and widely cultivated in the worldwide especially in southeastern Europe, northern Africa, and

northern Iran at the south end of the Caspian Sea [1]. Mediterranean region is the major production area of olive and there is a strong association with the high level of olive consumption and its associated products in this

Corresponding Author : Farshad H. Shirazi, Faculty of Pharmacy, Shahid Beheshti University of Medical Sciences, Tehran, Iran, SBMU Pharmaceutical Research Center, Tehran, Iran
Tel:+9888209627

E-Mail: F.shirazi@sbmu.ac.ir

Cite this article as: Bahramiun H, Kamalinejad M, Elahian F, Noubarani M, Balaghi F, Keyhanfar F, Eskandari M, H. Shirazi f, The Role of *Olea Europaea* L. Fruit on A2780, A172 and HFFF2 Proliferation. Iranian Journal of Pharmaceutical Sciences, 2016, 12 (2): 55-62.

region and reduced incidence of cardiovascular diseases and cancers of the breast, skin, and colon [2, 3]. It has been also reported the low incidence of chronic degenerative disease in Mediterranean countries [4]. Olive is traditionally used as an herbal remedy for the treatment of aphrodisiacs, emollients, laxatives, nutritive, sedatives, tonics, colic, alopecia, paralysis, rheumatic pain, sciatica, and hypertension [5]. In the context of religious importance, olive is narrated over several times in the Bible as well as is praised as a blessed fruit in Chapter 24 Al-Nur Quran [6]. The fruit and compression extracted oil, olive oil, possess various nutritional and medicinal properties since ancient times. Olive fruit contains valuable nutrients and bioactive compounds such as proteins, flavonoids, oleic acid, and phenolic compounds [1, 7]. Oleuropein is one of the most abundant phenolic constituents of olive with a considerable antioxidant activity [8]. Several studies have demonstrated that olive possesses a broad spectrum of pharmacological and therapeutic effects such as anti-inflammatory [9] and anticancer [10]. hepatoprotective activity [11], cardiogenic [12], antiatherogenic [13], anticarcinogenic [14], antimicrobial [15],

antihypertensive [16], and antidyslipidemic [17]. In this investigation it was aimed to explore the role of aqueous extract of *Olea europaea* L. fruit on A2780, A172 and HFFF2 cell proliferation by MTT assay. A2780 and A172 are human ovarian cancer cell line and human glioblastoma cell line, respectively; while HFFF2 is a normal human embryonic cell line. In this way we can evaluate the effects of olive on cancerous and normal cells proliferation in a new point of view.

2. Material and Methods

2.1. Materials

Fetal bovine serum (FBS), phosphate-buffered saline (PBS), trypsin, penicillin, streptomycin, DMSO, 3-4, 5-dimethylthiazol-2-yl-2, 5-diphenyl tetrazolium bromide (MTT), and the RPMI-1640 medium supplemented with 10% heat inactivated FBS were purchased from Autocell. The cell lines were provided by Pasteur Institute of Iran.

2.2. Plant Materials and the Preparation of the Extracts

Olea europaea L. fruit were collected from Tarom, Zanjan province of Iran. The collected fruits were scientifically approved by the Department of Botany, Shahid Beheshti University (Voucher number: 1115, deposited in: Shahid Beheshti University Herbarium). Fresh fruits were cleaned and then dried in the shade at room temperature. For the preparation of aqueous extract the fruits were decocted in water for 30 min. Then, the extract was filtered

and concentrated to the desired level (honey-like viscosity), and stored at -20°C. The moisture level of the extract was determined as follows: 2g of final extract was placed in an oven at 60–65°C for 72 h and then weighed. Weight loss was used as a moisture indicator. The final extract contained 24% water. These extracts were dissolved in distilled water at the desired concentrations just before use [18, 19].

2.3. Cell Culture and Cell Proliferation Assay By MTT

Activity effects of aqueous fruit extract of *Olea europaea* L. were assessed on human cancer cell lines (A2780 and A172) and human embryonic cell line (HFFF2) by using MTT (3-(4, 5-dimethylthiazol-2-yl)-2, 5-diphenyl tetrazolium bromide) assay as previously described [20, 21]. The cells were cultured in RPMI 1640 medium (A2780) or DMEM (A172 and HFF-2) enriched with 10% FBS (Fetal Bovine Serum) and incubated at 37°C with 5% CO₂ and 96% humidity. After several subcultures, cells were distributed in 96-well plates at 1,000 cells in 100 µL of culture medium and incubated for 24 h at same condition to allow attachment of cells to the bottom of wells. Then culture medium removed and 100 µL of ten-fold serially diluted concentrations of plant extracts (0.0001-100 mg/ml) added to each well in pentaplicate. Microtiter plates further incubated for 5 days in same condition. Culture medium without extract was used as negative control. After the

incubation time, the extract containing medium discharged and for evaluation of cell survival, 25 µL of MTT solution (4 mg/ml in PBS) added to each well and plates incubated for 3 h (in same condition). Then 100 µL of DMSO added to each well and plates were gently shaken to dissolve the formed formazan crystals. The absorbance of each well measured at 540 nm using an ELISA plate reader (Infinite M200, Tecan). The GI% (Growth Inhibition percent) was calculated using the formula %Growth Inhibition = $100 - (OD_{test} - OD_{control}) \times 100$, where OD_{test} is the mean absorbance of treated cells and OD_{control} is the mean absorbance of a negative control. The cell survival of control assumed 100% and cell growth values generated from dose-response curves for each cell line.

2.4. Statistical Analysis

All experiments were done in pentaplicate and the results were calculated as a mean ± standard deviation (SD). The experimental data were processed using the paired sample t-test and one way ANOVA analysis of the SPSS version 16.0 software for Windows.

3. Results and Discussion

In this study, human ovarian cancer cells (A2780) and human glioblastoma (A172) and also human embryonic cell line (HFFF2) were treated with different concentrations (0.0001-100 mg/ml) of aqueous extract of *Olea europaea* L. fruit. The cell lines were observed by an inverted Microscope and it was observed that




Figure 1. Determination of cell viability by an inverted microscope. The cell lines viability was detected by an inverted microscope (Axiovert 25 Inverted Microscope) just before MTT assay. A: A2780, B: A172, C: HFFF2.

they were viable during the study and before MTT assay (Figure 1). Cell viability was determined 5 days after the treatment. As indicated in figure 2, *Olea europaea* L. fruit extract increased A2780 proliferation at different concentration in comparison with control one (without extract). Also, the extract of *Olea europaea* L. induced the proliferation of A172 cell line at different concentration in comparison to control after the exposure (Figure 3). Again, as shown in figure 4, olive aqueous extract exhibited activatory effects on normal cell line HFFF2 proliferation at different concentration. Interestingly, *Olea europaea* L. extract had more activatory effect on A2780 in comparison with other two cell lines (Figure 5).

The olive tree, *Olea europaea* L., is a small evergreen tree that cultivated in different part of the world especially in Mediterranean [22, 23]. Olives and their constituents show important roles in diseases prevention through different mechanisms such as: anti-oxidant, anti-inflammatory, hepatoprotective activity, cardiotoxic activity. It is believed that olive

exerts its biological effects mainly via constituent antioxidants, including oleic acid, phenolics, and squalene [24, 25]. It has been also reported that hydroxytyrosol is one of the main phenolic components of olive oil that is responsible for chemopreventive effects of olive and olive oil [14, 26]. In addition, this active compound showed a preventive role in cardiovascular diseases and acquired immunodeficiency syndrome (AIDS). It has been proven that these preventive effects are mediated by its potent antioxidant and anti-inflammatory properties [27].

As shown in figure 2-4, aqueous extract of *Olea europaea* L. increased cell proliferation in human ovarian cancer cells (A2780), human glioblastoma (A172) and human fetus derived cells (HFFF2) in a dose dependent manner. However, the previous reports have been shown that *Olea europaea* L. fruit has antimutagenic effects [10]. These differences can be explained by different properties of cell lines, different types of date fruits and constitutes that fractionated in different solvents.




Figure 2. Effects of aqueous extract of *Olea europaea* L. fruit on cell proliferation of A2780. MMT assay was used for cell viability detection. Values are presented as mean±SD. *** $P < 0.001$ compared with control.




Figure 3. Effects of aqueous extract of *Olea europaea* L. fruit on cell proliferation of A172. MMT assay was used for cell viability detection. Values are presented as mean±SD. ** $P < 0.01$ compared with control.

In addition, the amounts of anticancer compounds that are responsible for antitumor effects of olive may be low in olives of our study. We believe that further investigation is needed to clarify the exact role of *Olea*

europaea L. in cell proliferation and cancer prevention. In this study the extract of *Olea europaea* L. showed more activatory effects on A2780 cell line in comparison with A172 and HFFF2. These differences in the activatory




Figure 4. Effects of aqueous extract of *Olea europaea* L. fruit on cell proliferation of HFFF2. MMT assay was used for cell viability detection. Values are presented as mean±SD. *** $P < 0.001$ compared with control.




Figure 5. Effects of *Olea europaea* L. aqueous extract on cell proliferation of A2780, A172, and HFFF2. MMT assay was used for cell viability detection. Values are presented as mean±SD. ** $P < 0.01$, *** $P < 0.001$ compared with HFFF2; 00 $P < 0.01$; 000 $P < 0.001$ compared with A172.

effects may be related to the activation of different signaling pathways in different cell lines.

4. Conclusion

In this study, the activatory effects of aqueous extract of *Olea europaea* L. were investigated in different cancerous (A2780 and

A172) and fetus derived cell line (HFFF2). The aqueous extract increased cell proliferation when exposed to these cell lines. On the basis of observed results further studies are necessary to confirm our findings. In the next steps the active constituents of aqueous extract should be isolated and tested for the confirmation and validation of our findings.

Acknowledgments

The results presented in this article were extracted from the thesis of Dr. Hadis Bahramiun (Pham.D. graduate of School of Pharmacy, Zanjan University of Medical Sciences) who performed her thesis under the supervision of Dr. Mohammad Reza Eskandari. This study was funded by a Discovery Grant from Zanjan University of Medical Sciences, Deputy of Research (Grant No: A-12-501-17).

References

[1] Ghanbari R, Anwar F, Alkharfy KM, Gilani AH, Saari N. Valuable Nutrients and Functional Bioactives in Different Parts of Olive (*Olea europaea* L.)-A Review. *Int J Mol Sci.* 2012; 13(3):3291–340.

[2] Keys A, Menotti A, Karvonen MJ, Aravanis C, Blackburn H, Buzina R, Djordjevic BS, Dontas AS, Fidanza F, Keys MH, et al. The diet and 15-year death rate in the Seven Countries Study. *Am J Epidemiol.* 1986; 124:903-15.

[3] Owen RW, Haubner R, Würtele G, Hull E, Spiegelhalter B, Bartsch H. Olives and olive oil in cancer prevention. *Eur J Cancer Prev.* 2004; 13:319-26.

[4] Harwood JL, Yaqoob P. Nutritional and health aspects of olive oil. *Eur J Lipid Sci Technol.* 2002; 104:685-97.

[5] Hashmi MA, Khan A, Hanif M, Farooq U, Perveen S. Traditional Uses, Phytochemistry, and Pharmacology of *Olea europaea* (Olive). *Evid Based Complement Alternat Med.* 2015; 2015: 541591

[6] Ryan D, Robards K. Phenolic compounds in olives. *Analyst.* 1998, 123:31R–44R.

[7] Covas MI, Nyyssönen K, Poulsen HE, Kaikkonen J, Zunft HJ, Kiesewetter H, Gaddi A, de la Torre R, Mursu J, Bäumler H, Nascetti S, Salonen JT, Fitó M, Virtanen J, Marrugat J; EUROLIVE Study Group. The effect of

polyphenols in olive oil on heart disease risk factors: a randomized trial. *Ann Intern Med.* 2006; 145(5):333-41.

[8] Owen RW, Giacosa A, Hull WE, Haubner R, Würtele G, Spiegelhalter B, Bartsch H. Olive oil consumption and health: the possible role of antioxidants. *Lancet Oncol.* 2000; 1:107-112.

[9] Sahranavard S, Kamalinejad M, Faizi M. Evaluation of Anti-Inflammatory and Anti-Nociceptive Effects of Defatted Fruit Extract of *Olea europaea*. *Iran J Pharm Res.* 2014; 13(Suppl):119-23.

[10] Hamdi HK, Castellon R. Oleuropein, a nontoxic olive iridoid, is an anti-tumor agent and cytoskeleton disruptor. *Biochem Biophys Res Commun.* 2005; 334:769-78.

[11] Domitrović R, Jakovac H, Marchesi VV, Šain I, Romić Ž, Rahelić D. Preventive and therapeutic effects of oleuropein against carbon tetrachloride-induced liver damage in mice. *Pharmacol Res.* 2012; 65(4): 451-64.

[12] de Lorgeril M, Salen P. The Mediterranean diet in secondary prevention of coronary heart disease. *Clin Invest Med.* 2006; 29:154-8.

[13] Aviram, M. Interaction of oxidized low density lipoprotein with macrophages in atherosclerosis, and the antiatherogenicity of antioxidants. *Eur J Clin Chem Clin Biochem.* 1996, 34:599–608.

[14] Fabiani R, De Bartolomeo A, Rosignoli P, Servili M, Montedoro GF, Morozzi G. Cancer chemoprevention by hydroxytyrosol isolated from virgin olive oil through G1 cell cycle arrest and apoptosis. *Eur J Cancer Prev.* 2002; 11:351-8.

[15] Romero C, Medina E, Vargas J, Brenes M, De Castro A. In vitro activity of olive oil polyphenols against *Helicobacter pylori*. *J Agric Food Chem.* 2007; 55:680-6.

[16] Carollo C, Presti RL, Caimi G. Wine, diet, and arterial hypertension. *Angiology.* 2007; 58:92-6.

[17] Fitó M, Covas MI, Lamuela-Raventós RM, Vila J, Torrents L, de la Torre C, Marrugat J. Protective effect

of olive oil and its phenolic compounds against low density lipoprotein oxidation. *Lipids*. 2000; 35:633–8.

[18] Mirmohammadlu M, Hosseini SH, Esmaeili Gavgani M, Kamalinejad M, Noubarani M, Eskandari MR. Hypolipidemic, hepatoprotective and renoprotective effects of *Cydonia oblonga* Mill. fruit in streptozotocin-induced diabetic rats. *Iran J Pharm Res*. 2015;14(4): 1207-14.

[19] Heidari H, Kamalinejad M, Noubarani M, Rahmati M, Jafarian I, Adiban H, Eskandari MR. Protective mechanisms of *Cucumis sativus* in diabetes-related model of oxidative stress and carbonyl stress. *Bioimpacts*. 2016; 6(1):33-9.

[20] Jamali B, Nakhjavani M, Hosseinzadeh L, Amidi S, Nikounezhad N, Shirazi F. Intracellular GSH Alterations and Its Relationship to Level of Resistance following Exposure to Cisplatin in Cancer Cells. *Iran J Pharm Res*. 2015; 14(2):513-9.

[21] Ramazani A, Mashhadi M, Kamalinejad M, Tavakolizadeh M, Balaghi F, Noubarani M, Eskandari MR, Shirazi F.H. *Phoenix dactylifera* L.

Fruit induces cell proliferation of A2780, A172 and HFFF2 cell lines. *Iran J Pharm Res*. 2016; 12(1): 35-44.

[22] Fares R, Bazzi S, Baydoun SE, Abdel-Massih RM. The antioxidant and anti-proliferative activity of the Lebanese *Olea europaea* extract. *Plant Foods Hum Nutr*. 2011; 66(1):58-63.

[23] Turkez H, Togar B, Polat E. Olive leaf extract modulates permethrin induced genetic and oxidative damage in rats. *Cytotechnology*. 2012; 64(4):459–64.

[24] Waterman E, Lockwood B. Active components and clinical applications of olive oil. *Altern Med Rev*. 2007; 12(4):331-42.

[25] Tuck KL, Hayball PJ. Major phenolic compounds in olive oil: metabolism and health effects. *J Nutr Biochem*. 2002; 13:636-44.

[26] Visioli F, Poli A, Gall C. Antioxidant and other biological activities of phenols from olives and olive oil. *Med Res Rev*. 2002; 22:65-75.

[27] Vilaplana-Pérez C, Auñón D, García-Flores LA, Gil-Izquierdo A. Hydroxytyrosol and Potential Uses in Cardiovascular Diseases, Cancer, and AIDS. *Front Nutr*. 2014; 1: 18.